

Ankle Posterior Clearance

What is the problem?

Posterior ankle impingement is pain at the back of the ankle generally caused by a prominent piece of bone.

What's involved?

Posterior clearance surgery involves:

- general anaesthetic, intravenous antibiotics
- incision at the back of the ankle
- dissection down to the bony prominence (os trigonum)
- removal of the bone and tissue causing impingement
- wound closure with stitches/sutures
- infiltration of local anaesthetic

What's the worst thing(s) that can happen with surgery?

All surgical procedures carry some risk. Some of the risks of posterior clearance surgery include:

- Infection
- Problems with wound healing
- Nerve injury causing numbness, tingling and/or pins and needles
- Ankle stiffness
- Deep venous thrombosis/pulmonary embolism. (The risk of DVT increase with smoking, the oral contraceptive pill and hormone replacement therapy, immobility and obesity)
- Anaesthetic complications
- Drug allergy
- Ongoing pain

What can I expect after surgery?

Rest

- You will have a boot on your leg. Keep foot elevated as much as possible, especially for initial 72 hours. Keep dressings dry and intact until post operative appointment.
- Mobilise short distances as pain allows, aim to remain in your own home for the first few weeks limiting long periods of standing or mobilising

- Bear weight as tolerated in the boot

Pain

- Pain relief may be required for up to 2 weeks; depending on your other medications this may include:
 - Regular paracetamol, anti-inflammatory drugs and break through opiate type medications

Driving

- Any surgery performed on the right foot or both feet will mean no driving for a minimum of 2-6 weeks
- If surgery was performed on the left foot then driving in an automatic is possible, but Dr Phegan would recommend waiting a minimum of two weeks before attempting to drive

Driving any vehicle with a cast or boot is potentially hazardous and you should consider this before getting behind the wheel

Follow up

- Keep all dressings dry and in tact for the first two weeks
- Removal of stitches/sutures typically occurs at an appointment with Dr Phegan and his in-house therapists at 10-14 days at first post operative appointment. Very occasionally some sutures need to stay in for longer
- Your appointment will be booked by Dr Phegan and a confirmatory SMS will be sent informing you of the time and date for follow up

Typical milestones

- 2 weeks: Wound review and removal of sutures
- 2-6 weeks: regular physiotherapy to increase range and weight bearing
- 6weeks to 3 months: work on swelling reduction, scar massage and mobility
- Full recovery 3-6 months

Every patient's recovery is individual and depends on the severity of the injury and the complexity of the surgery.

ANY PROBLEMS

During office hours contact Dr Phegan's office on (07) 56711120 or email info@drphegan.com

After hours please contact the hospital where your surgery was performed or attend your local GP